

*Main panel*

---

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

**Snailex**  
**Slug and Snail Pellets**

**ACTIVE CONSTITUENT: 15 g/kg METALDEHYDE**

**For the control of slugs and snails in many agricultural and horticultural situations as specified in the Directions for Use table.**

**Net Contents: (20 – 1,000 kg, 22,000 kg)**

**Agribait Pty Ltd**

ABN 55 162 181 884

**Lot 33/46 Pine Freezers Road, Port Lincoln SA 5606**

**Phone: 08 7626 8002**

**APVMA APPROVAL NO.: 68580/143036**

**DIRECTIONS FOR USE**

**Restraint:** DO NOT place pellets in heaps.

SITUATION	PEST	RATE	CRITICAL COMMENTS
			Use the higher rate where there are large populations of snails or slugs and where wet cool climatic conditions are favourable for snail and slug movement.
Agricultural crops, cereals, oilseeds (including poppies) Horticultural crops Fruit (including grapevines, vegetables)	Snails and slugs including: Common white snail ( <i>Ceruella virgata</i> ) White Italian snail ( <i>Theba pisana</i> ) Conical snail ( <i>Cochlicella acuta</i> ) Grey field slug ( <i>Deroceras reticulatum</i> )	5-10 kg/ha	Broadcast pellets evenly over the surface of the ground where snails and slugs are active. Brown field slug mortality may be delayed, requiring longer feeding times.
Pulses, Pasture establishment	Brown field slug ( <i>Deroceras invadens</i> ) (excludes Small Brown Snails)		This product is to be used only during the establishment stages of pulse crops and pastures. Broadcast pellets evenly over the crop area to be protected either prior to planting, post-plant pre-emergence or early post-emergence (prior to 4 leaf stage).
Other agricultural, industrial and commercial situations where snails and slugs congregate			Broadcast pellets evenly over the surface of the ground where snails and slugs are active.
Around plants. Around crops and/or border areas where snails and slugs congregate.	Slugs, Snails	50 g/m <sup>2</sup>	Scatter the pellets around the area to be protected and/or where snails and slugs congregate. Spread pellets evenly.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**HARVEST**

**Fruit and Vegetables: DO NOT APPLY LATER THAN 7 DAYS BEFORE HARVEST.**

**Pulses, Oilseeds, Winter cereals: NOT REQUIRED WHEN USED AS DIRECTED.**

**GRAZING**

**Pastures, Pulses, Oilseeds, Winter cereals: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 5 WEEKS AFTER APPLICATION.**

## **GENERAL INSTRUCTIONS**

Scatter pellets using an appropriate applicator. The best time to apply is immediately after rain or during humid overcast weather when slugs and snails are most active. The pellets eventually decompose leaving no harmful residue. Use regularly to control re-infestation.

## **TIMING**

The best time to apply is immediately after rain, or during humid overcast weather when slugs and snails are most active. For broadacre crops, start when rain/moisture triggers snail activity in Autumn and before significant egg laying.

## **PRECAUTIONS**

DO NOT apply to edible plant parts. DO NOT graze or cut for stockfeed any crops while pellets are visible in the plant foliage.

## **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers.

## **PROTECTION OF DOMESTIC ANIMALS AND LIVESTOCK**

Keep away from animals, poultry, ducks and domestic pets as accidental poisoning may be fatal. Dogs and other animals find this bait attractive and it may kill them. If pets or livestock are poisoned contact a veterinary surgeon.

## **STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well-ventilated area, out of direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. Shake and empty into applicator. DO NOT dispose of unused product on-site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. DO NOT burn empty containers or product.

## **SAFETY DIRECTIONS**

Poisonous if swallowed. Wash hands after use.

## **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 13 11 26, New Zealand 0800 764 766.

## **SAFETY DATA SHEET**

Additional information is listed in the safety data sheet (SDS). A copy is available from the supplier on request.

## **CONDITIONS OF SALE:**

The use of this product being beyond the control of the manufacturer, no warranty expressed or implied is given by the manufacturer regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and the manufacturer accepts no responsibility for any consequence whatsoever resulting from the use of this product.

Batch No.

Date of manufacture: